

CM 22027 - RAPPAPORT VASSILIADIS SALMONELLA ENRICHMENT BROTH (as per USP/BP/EP/JP/IP)

INTENDED USE

For selective enrichment & isolation of *Salmonella* species from pharmaceutical products.

PRODUCT SUMMARY AND EXPLANATION

Rappaport Vassiliadis Salmonella Enrichment Broth is used for the enrichment and selective isolation of *Salmonella* species. It conforms to harmonized USP/EP/JP/BP/IP requirements. Rappaport et al. formulated an enrichment medium for *Salmonella* species. Which was modified by Vassiliadis et al. Rappaport Vassiliadis Salmonella Enrichment Broth medium is evaluated as an alternative of Rappaport-Vassiliadis (RV) Broth where Enzymatic Digest of Casein is replaced by Soya Peptone as the nitrogen and vitamin source.

COMPOSITION

Ingredients	Gms / Ltr
Magnesium chloride hexahydrate	29.000
Sodium chloride	8.000
Soya peptone	4.500
Potassium dihydrogen phosphate	0.600
Dipotassium phosphate	0.400
Malachite green	0.036

PRINCIPLE

The medium contains Sodium chloride which maintains the osmotic balance in the medium. The low pH of the medium, combined with the presence of Malachite green and Magnesium chloride raises the osmotic pressure and provides selectivity for the highly resistant *Salmonella* species. Potassium dihydrogen phosphate and dipotassium phosphate buffers the medium to maintain a constant pH. Soya peptone acts as a source of essential nutrients & vitamins for enhancing the growth of *Salmonella*.

INSTRUCTION FOR USE

- Dissolve 42.50 grams of the medium in 1000 ml distilled water.
- Gently heat to boiling with swirling to dissolve the medium completely.
- Dispense as desired into tubes.
- Sterilize by autoclaving at 15 psi (115°C) for 30 minutes or as per validated cycle
- Cool to room temperature.

QUALITY CONTROL SPECIFICATIONS

Appearance of Dehydrated powder	:	Light yellow to light blue, homogeneous free flowing powder
Appearance of Prepared medium	:	Greenish blue coloured, clear to slightly opalescent solution
pH (at 25°C)	:	5.2±0.2



INTERPRETATION

Culture characteristics observed after incubation. Recovery was carried out using XLD Agar after enrichment in Rappaport Vassiliadis Salmonella Enrichment Broth.

Microorganism	ATCC	Inoculum (CFU/ml)	Growth	Colour of colony on XLD	Recovery	Incubation Temperature	Incubation Period
<i>Salmonella typhimurium</i>	14028	50-100	Luxuriant	Red with black centres	≥70%	30-35°C.	18-24 Hours
<i>Salmonella enteritidis</i>	13076	50-100	Luxuriant	Red with black centres	≥70%	30-35°C.	18-24 Hours
<i>Salmonella arizonae</i>	13314	50-100	Luxuriant	Red with black centres	≥70%	30-35°C.	18-24 Hours
<i>Escherichia coli</i>	25922	50-100	None-Poor	Yellow	0-10%	30-35°C.	18-24 Hours
<i>Escherichia coli</i>	8739	50-100	None-Poor	Yellow	0-10%	30-35°C.	18-24 Hours
<i>Staphylococcus aureus</i>	25923	≥1000	Inhibited	-	0%	30-35°C.	18-24 Hours
<i>Staphylococcus aureus</i>	6538	≥1000	Inhibited	-	0%	30-35°C.	18-24 Hours

PACKAGING

In 100 & 500 gm packaging size.

STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers below 25°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out, replace the cap tightly to protect from hydration.

Product Deterioration: Do not use, if powder shows evidence of microbial contamination, discoloration, drying, or other signs of deterioration.










DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

REFERENCES

- The United States Pharmacopoeia. Amended Chapters 61, 62 & 111, The United States Pharmacopoeial Convention Inc., Rockville, MD. (2009).
- Directorate for the Quality of Medicines of the Council of Europe (EDQM). The European Pharmacopoeia, Amended Chapters 2.6.12, 2.6.13, 5.1.4, Council of Europe, 67075 Strasbourg Cedex, France. (2007).
- Japanese Pharmacopoeia. Society of Japanese Pharmacopoeia. Amended Chapters 35.1, 35.2, 7. The Minister of Health, Labor, and Welfare. (2008).
- Rappaport, F., N. Konforti, and B. Navon. A new enrichment medium for certain salmonellae. J. Clin. Pathol. 9:261-266. (1956).
- Vassiliadis, P., D. Trichopoulos, A. Kalandidi, and E. Xirouchaki. Isolation of salmonellae from sewage with a new procedure of enrichment. J. Appl. Bacteriol. 44:233-239. (1978).
- Van Schothorst, M. and A. M. Renaud. J. Appl. Bact. 54:209-215. (1983).
- McGibb
- on, L., E. Quail, and C. R. Fricker. Inter. J. Food Microbiol. 1:171-177. (1984).



 GMP Good Manufacturing Practices Certified	 Best Before	 QTY. Quantity	 REF Catalogue Number	 Manufacturer
 Temperature Unit	 LOT/ B. NO. Lot / Batch Number	 Consults Instructions for Use	 QR Code	

NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

***For professional use only.**

