

## CM 22056 – EMB AGAR, LEVINE

### INTENDED USE

For isolation, enumeration and differentiation of members of Enterobacteriaceae from pharma, dairy & food products.

### PRODUCT SUMMARY AND EXPLANATION

Levine EMB Agar was developed by Levine and is used for the differentiation of Escherichia coli and Enterobacter aerogenes and also for the rapid identification of Candida albicans. This medium is recommended for the detection, enumeration and differentiation of members of the coliform group by American Public Health Association. It is also recommended by BIS for detection and estimation of coliform bacteria in food stuff and Escherichia coli from food and water.

### COMPOSITION

Ingredients	Gms / Ltr
Peptic digest of animal tissues	10.00
Dipotassium phosphate	2.000
Lactose	10.00
Eosin -Y	0.400
Methylene blue	0.065
Agar	15.000

### PRINCIPLE

Eosin-Y and methylene blue make the medium slightly selective and inhibit certain gram-positive bacteria. These dyes differentiate between lactose fermenters and non-fermenters. Some gram-positive bacteria such as faecal Streptococci, yeasts grow on this medium and form pinpoint colonies. Weld proposed the use of Levine EMB Agar, with added Chlortetracycline hydrochloride, for the rapid identification of Candida albicans in clinical specimens. A positive identification of Candida albicans can be made after 24 - 48 hours' incubation at 35 - 37°C in 10% carbon dioxide atmosphere, from specimens such as faeces, oral and vaginal secretions and nail or skin scraping etc. However, the typical appearance is variable.

### INSTRUCTION FOR USE

Dissolve the 37.5 grams in 1000 ml distilled water.

Heat to boiling to dissolve the medium completely.

Sterilize by autoclaving at 15 psi pressure (121°C) for 15 minutes. Avoid overheating.

Cool to 50°C and shake the medium in order to oxidize the methylene blue (i.e. restore its blue colour) and to suspend the precipitate which is an essential part of the medium.

Precaution: Store the medium away from light to avoid photo oxidation.

### QUALITY CONTROL SPECIFICATIONS



Appearance of Powder : Light pink to purple coloured homogeneous free flowing powder.  
 Appearance of prepared medium : Reddish purple coloured slightly opalescent gel with greenish cast and finely dispersed precipitate, forms in Petri plates.  
 pH (at 25°C) : 7.1 ± 0.2

#### INTERPRETATION

Cultural characteristics observed after an incubation.

Microorganism	ATCC	Inoculum (CFU/ml)	Growth	Recovery	Color of the colony	Incubation Temperature	Incubation Period
Candida albicans	10231	10-100	Good-Luxuriant (Incubated in 10% carbon dioxide)	>=70%	Colourless	35-37°C	24-48 Hours
Enterobacter aerogenes	13048	50-100	Good	40-50%	Pink red	35-37°C	24-48 Hours
Enterococcus faecalis	29212	>=10 <sup>3</sup>	Inhibited	0%	-	35-37°C	24-48 Hours
Escherichia coli	25922	50-100	Luxuriant	>=70%	Blue- black with metallic sheen	35-37°C	24-48 Hours
Pseudomonas aeruginosa	27853	50-100	Luxuriant	>=70%	Colourless	35-37°C	24-48 Hours
Saccharomyces cerevisiae	9763	10-100	None-poor	0-10%	Cream	35-37°C	24-48 Hours
Salmonella Serotype Typhimurium	14028	50-100	Luxuriant	>=70%	Colourless	35-37°C	24-48 Hours
Staphylococcus aureus	25923	50-100	None-poor	0-10%	Colourless	35-37°C	24-48 Hours

#### PACKAGING:

In pack size of 100 gm and 500 gm bottles.



## STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.


Product Deterioration: Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.

## DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

## REFERENCES

1. Levine M., 1918, J. Infect. Dis., 23:43.
2. Levine M., 1921, Bull. 62, Iowa State College Engr. Exp. Station.
3. Greenberg A. E., Trussell R. R. and Clesceri L. S. (Eds.), 1985, Standard Methods for the Examination of Water and Waste water, 16th ed., APHA, Washington, D.C.
4. Marshall R. (Ed.), 1992, Standard Methods for the Examination of Dairy Products, 16th ed., APHA Inc., New York.
5. Speck M. (Ed.), 1992, Compendium of Methods for the Microbiological Examination of Foods, 3rd ed., APHA, Washington, D.C.
6. Bureau of Indian Standards, IS : 5401, 1969 (Second reprint - June 1990).
7. Bureau of Indian Standards, IS : 5887 (Part - I) 1976, reaffirmed 1986.
8. Weld J. T., 1952, Arch. Dermat. Syph., 66:691.
9. Weld J. T., 1953, Arch. Dermat. Syph., 67(5):433.



NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

\*For LabUse Only

