

CM 22245 - CHROMOGENIC COLIFORM AGAR PLATE

INTENDED USE

For determination of coliform bacteria particularly Enterobacteriaceae on the basis of their ability to ferment lactose.

PRODUCT SUMMARY AND EXPLANATION

Coliforms including *Escherichia coli* are used as primary indicators of faecal contamination in water and food industries. Their presence and enumeration in samples is used as an index of the presence of faecal matter and is indicative of the possible presence of enteric pathogens. Chromogenic Coliform Agar (CCA) is a fast, accurate, and efficient way to detect coliforms bacteria particularly Enterobacteriaceae during microbiological quality testing of water and food samples. *Escherichia coli* are the most common pathogen in urinary tract infections. This product incorporates the company's chromogenic galactoside that detects clinical levels of coliforms with high sensitivity.

COMPOSITION

Ingredients	Gms / Ltr
Agar	10.00
Sodium Chloride	5.00
Disodium hydrogen phosphate	2.70
Sodium dihydrogen phosphate (2H ₂ O)	2.20
Yeast extract	2.00
Enzymatic digest of casein	1.00
Sodium pyruvate	1.00
Sorbitol	1.00
Tryptophan	1.00
Salmon-β-D-galactoside	0.20
Sodium heptadecylsulphate (Tergitol 7)	0.15
X-glucuronide	0.10
Isopropyl 1-β-D- thiogalactopyranoside (IPTG)	0.10

PRINCIPLE

This medium contains enzymatic digest of casein, yeast extract, sorbitol and sodium pyruvate as sources of carbon, nitrogen, fermentable carbohydrate and other essential growth nutrients for the growth of microorganisms. Disodium hydrogen phosphate and sodium dihydrogen phosphate are the buffering agents. Sodium chloride maintains the osmotic equilibrium in the medium. L- Tryptophan improves the indole reaction and gives improved differentiation between *Escherichia coli* and other coliforms. Tergitol- 7 inhibits gram positive bacteria. The two chromogens used; salmon-β-D-galactoside and X-glucuronide helps in differentiation of *Escherichia coli* and other coliforms on the basis of colony colour. The enzyme β-D-galactosidase cleaves salmon-β-D-galactoside, and gives a salmon to red colour to the coliform bacteria. *E. coli* have β-D galactosidase and β-D-glucuronidase enzymes to cleave both the chromogens, which give blue-violet colour to colonies. Expression of β-D galactosidase is strengthened in the presence of IPTG. Agar is a gelling agent.

INSTRUCTION FOR USE

Either streak, inoculate or surface spread the test inoculum aseptically on the plate.



QUALITY CONTROL SPECIFICATIONS

Appearance	:	Light yellow coloured, clear to slightly opalescent gel.
Quantity of Medium	:	25ml of medium in 90mm plates.
pH (at 25°C)	:	6.8 ± 0.2
Sterility Check	:	Passes release criteria

INTERPRETATION

Cultural characteristics observed after incubation.

Microorganism	ATCC	Inoculum (CFU/ml)	Growth	Recovery	Colour of Colony	Incubation Temperature	Incubation Period
<i>Escherichia coli</i>	25922	50-100	Luxuriant	>=70%	Dark blue to violet	35-37°C	18-24 Hours
* <i>Klebsiella aerogenes</i>	13048	50-100	Luxuriant	>=70%	Pink to red	35-37°C	18-24 Hours
<i>Citrobacter freundii</i>	43864	50-100	Luxuriant	>=70%	Pink to red	35-37°C	18-24 Hours
<i>Enterococcus faecalis</i>	19433	≥1000	Inhibited	0%	-	35-37°C	18-24 Hours
<i>Salmonella</i> Enteridis	13076	50-100	Luxuriant	>=70%	Colourless	35-37°C	18-24 Hours

*Formerly known as *Enterobacter aerogenes*

PACKAGING:

Doublelayered packing containing 5 No. of plates with one silica gel desiccant bag packed inside it.

STORAGE

On receipt, store the plates at 2–8 °C. Avoid freezing and overheating. Do not open until ready to use. Prepared plates stored in their original sleeve wrapping until just prior to use may be inoculated up to the expiration date and incubated for recommended incubation times. Allow the medium to warm to room temperature before inoculation.











Product Deterioration: Do not use plates if they show evidence of microbial contamination, discoloration, drying, cracking or other signs of deterioration.

DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

REFERENCES

- ISO9308-1/2014. Water quality — Enumeration of *Escherichia coli* and coliform bacteria —Part 1: Membrane filtration method for waters with low bacterial background flora.
- Frampton, E.W., Restaino, L. and Blaszkowski, N. 1988. J. Food Prot. 51: 402-404.
- Kilian, M. and Bulow, P. 1979. Acta. Pathol. Microbiol. Scand. (Section B) 87: 271-276.

 Quantity	 Lot / Batch Number	 Temperature Unit	 Best Before	 Manufacturer
 Certification of Good Manufacturing Practices	 Catalogue No.	 European Conformity	 QR Code	 Consults Instructions for use :



NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

***For Lab Use Only**

