

CM 20322 – BRYANT AND BURKEY AGAR

INTENDED USE

For detection and enumeration of spores of lactate fermenting Clostridium in dairy products.

PRODUCT SUMMARY AND EXPLANATION

Bryant and Burkey Agar is used to enumerate the spores of gasogenic clostridia that are responsible for swelling of cheese in dairy industry. In normal conditions of use, the medium allows the growth of other microorganisms also which are not directly related to cheese alteration, e.g. Clostridium sporogenes or Clostridium butyricum. Germination and growth of butyric acid bacteria (BAB) causes severe defects in cheese with silage being the main source of BAB spores in cheese milk. Clostridia spores are heat resistant and therefore can contaminate cheese brines. The gas produced by the growth of clostridia swells the cheese and is responsible for defect known as butyric swelling, resulting in bad taste. The main species causing this butyric swelling defect is Clostridium tyrobutyrium.

Bryant and Burkey Agar is the modification of Bryant and Burkey Medium with addition of little amount of agar that makes the medium thicker to be used in greater volumes. Addition of sodium thioglycollate maintains anaerobic conditions and also the amount of lactate is reduced so that the density of the medium will be retained easily for the production of gas bubbles.

COMPOSITION

Ingredients	Gms / Ltr
Casein enzymic hydrolysate	15.000
Yeast extract	5.000
Beef extract	7.500
Sodium acetate	5.000
L-Cystine hydrochloride	0.600
Resazurin	0.0025
Sodium thioglycollate	0.200
Agar	0.750

PRINCIPLE

Resazurin is a redox indicator and monitors the oxygen level turning from pink under aerobic conditions to colourless under anaerobic conditions. The nutrient composition of the basal medium, particularly of casein enzymic hydrolysate, yeast extract, beef extract and L-cystine HCl help in rapid growth of Clostridium species. Sodium acetate promotes spore germination and improves the selectivity of the medium. Sodium lactate is fermented under anaerobic conditions by Clostridium tyrobutyricum. A pink colour indicates the presence of oxygen in the media which turns colourless on boiling.

INSTRUCTION FOR USE

Dissolve 34.05 grams in 1000 ml distilled water containing 3 grams of Sodium lactate.

Heat to boiling to dissolve the medium completely.

Sterilize by autoclaving at 15 psi pressure (121°C) for 15 minutes.

Note: If more than the upper one-third of the medium has acquired a pink colour, the medium may be restored once by heating in a water bath or in free flowing steam until the pink colour disappears.

QUALITY CONTROL SPECIFICATIONS



Appearance of Powder : Cream to yellow homogeneous free flowing powder.
 Appearance of prepared medium : Light amber coloured, clear to slightly opalescent solution with upper 10% or less medium pink on standing.
 pH (at 25°C) : 5.9±0.2

INTERPRETATION

Cultural characteristics observed after incubation.

Microorganism	ATCC	Inoculum (CFU/ml)	Growth	Gas production	Incubation Temperature	Incubation Period
Clostridium tyrobutyricum	25755	50-100	Good	Positive reaction	35-37°C	6 Days

PACKAGING:

In pack size of 500 gm bottles.

STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.










Product Deterioration: Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.

DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

REFERENCES

1. Bryant M. P. and Burkey L. A., 1956, J. Bacteriol., 71: 43-46.
2. Bergeres J. L. and Sivela S., 1989, Detection and enumeration of Clostridial spores related to cheese quality-Classical and new methods, FIL-IDF Bull. 51:18-23.
3. Touraille C. and Bergere J. L., 1974, Biochimie, 56: 404-422.
4. Vissers M. M. M., Drichuis F., Giffel T., John P. D., Lankveld J. M. G., 2007, J. Dairy Sci., 90:928

 GMP Good Manufacturing Practices Certified	 Best Before	 QTY. Quantity	 REF Catalogue Number	 Manufacturer
 Temperature Unit	 LOT/ B. NO. Lot / Batch Number	 Consults Instructions for Use	 QR Code	

NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

*For Lab Use Only

