

CM 23705 – ACRIFLAVIN-CEFSULODIN-VANCOMYCIN SUPPLEMENT (ACV SUPPLEMENT)

INTENDED USE

For the isolation of Enterohemorrhagic E. coli (EHEC).

COMPOSITION

Ingredients	Concentration
Acriflavin	50mg
Cefsulodin	40mg
Vancomycin	50mg

(Per vial sufficient for 500ml medium)

APPLIED IN MEDIUM






BUFFERED PEPTONE WATER W/ PYRUVATE

INSTRUCTION FOR USE

Rehydrate the contents of 1 vial aseptically with 5 ml of sterile distilled water. Mix well and aseptically add to 500 ml of sterile, cooled (45-50°C) Buffered peptone water w/ pyruvate. Mix well and dispense as desired.

STORAGE:

Vials should be stored in sealed container at 2-8°C.

 GMP Good Manufacturing Practices Certified	 IVD For In Vitro Diagnostic Use	 QTY. Quantity	 LOT/ B. NO. Lot / Batch Number	 REF Catalogue Number	 Manufacturer
 Temperature Unit	 EC REP Authorized Representative <small>MedWor GmbH Buckhorn 16 48147 Aachen, Germany</small>	 European Conformity	 QR Code	 Consults Instructions for Use	 Best Before

NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

*For Lab Use Only

