

CM 23828 – LISTERIA SELECTIVE SUPPLEMENT-II

INTENDED USE

For the selective isolation and cultivation of *Listeria monocytogenes*.

COMPOSITION

Ingredients	Concentration
Acriflavin hydrochloride (Trypaflavin)	5mg
Nalidixic acid	20mg
Cycloheximide	25mg

(Per vial sufficient for 500 ml medium)

INSTRUCTION FOR USE

Rehydrate the contents of one vial with 5 ml sterile distilled water. Mix well and aseptically add to 500 ml sterile, cooled (45-50°C) BUFFERED LISTERIA ENRICHMENT BROTH BASE (BLE Broth Base) (as per FDA) / LISTERIA SELECTIVE BROTH BASE / LISTERIA SELECTIVE BROTH BASE (VEG.) / LISTERIA SELECTIVE AGAR BASE. Mix well and dispense as desired.

STORAGE

Vials should be stored in sealed container at 2-8°C.

 Good Manufacturing Practices Certified	 For In Vitro Diagnostic Use	 Quantity	 Lot / Batch Number	 Catalogue Number	 Manufacturer
 Temperature Unit	 Authorized Representative MedWet GmbH Buckhorn 16, 48147 Münster, Germany	 European Conformity	 QR Code	 Consults Instructions for Use	 Best Before

NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

*For Lab Use Only

