

CM 21183 - SOYA CASEIN DIGEST AGAR W/ TWEEN 80 AND LECTHIN (MICROBIAL CONTENT TEST AGAR)

INTENDED USE

For detection and enumeration of microorganisms present on the surfaces of sanitary importance.

PRODUCT SUMMARY AND EXPLANATION

Tryptone Soya Agar with Lecithin and Polysorbate 80 is used in RODAC (Replicate Organism Detection and Counting) plates for the detection and enumeration of microorganisms present on surfaces of sanitary importances. Collection of samples from areas before and after the treatment with disinfectant evaluates cleaning procedures in environmental sanitation. The presence and number of microorganisms is determined by the appearance of colonies on the agar surface. After counting the colonies, carry out biochemical testing for identification.

COMPOSITION

Ingredients	Gms / Ltr
Casein enzymic hydrolysate	15.000
Soya peptone	5.000
Sodium chloride	5.000
Lecithin	0.700
Polysorbate 80 (Tween 80)	5.000
Agar	15.000

PRINCIPLE

Casein enzymic hydrolysate and soya peptone provide nitrogenous compounds and other nutrients essential for microbial replication. Lecithin and polysorbate 80 (Tween 80) are neutralizers reported to inactivate residual disinfectants from where the sample is collected. Lecithin neutralizes quaternary ammonium compounds and polysorbate 80 neutralizes phenolic disinfectants, hexachlorophene, formalin and with lecithin ethanol.

INSTRUCTION FOR USE

- Dissolve 45.7 grams in 1000 ml distilled water.
- Heat to boiling to dissolve the medium completely.
- Sterilize by autoclaving at 12 to 15 psi pressure (118 - 121°C) for 15 minutes.
- Cool to 45-50°C. Mix well and pour into sterile Petri plates.

QUALITY CONTROL SPECIFICATIONS

- Appearance of Powder : Cream to yellow coloured granular medium.
- Appearance of prepared medium : Light yellow to medium amber coloured clear to slightly opalescent gel forms in Petri plates.
- pH (at 25°C) : 7.3±0.2

INTERPRETATION

Cultural characteristics observed after an incubation.



Microorganism	ATCC	Inoculum (CFU/ml)	Growth	Recovery	Colour of colony	Incubation Temperature	Incubation Period
Staphylococcus aureus	25923	50-100	Luxuriant	$\geq 70\%$	Yellow to golden	35 - 37°C	18-24 Hours
Pseudomonas aeruginosa	27853	50-100	Luxuriant	$\geq 70\%$	Yellow green	35 - 37°C	18-24 Hours

PACKAGING:

In pack size of 100 gm and 500 gm bottles.

STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.

Product Deterioration: Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.

DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

REFERENCES

- Halland Hartnett, 1964, Public Hlth. Rep., 79:1021.
- Richardson (Ed.), 1985, Standard Methods for the Examination of Dairy Products, 15th ed., APHA, Washington, D.C.
- MacFaddin J.F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. I, Williams and Wilkins, Baltimore.
- Brummer, 1976, Appl. Environ. Microbiol., 32:80.
- Favero (Chairman), 1967, Biological Contamination Control Committee, a state of the art report., Am. Assoc. for contamination control.
- Murray P. R, Baron E. J., Jorgensen J. H., Tenover F. C., Tenover F. C., (Eds.), 2007, Manual of Clinical Microbiology, 9th Ed., ASM, Washington, D.C.

 Good Manufacturing Practices Certified	 Best Before	 Quantity	 Catalogue Number	 Manufacturer
 Temperature Unit	 Lot / Batch Number	 Consults Instructions for Use	 QR Code	

NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

*For Lab Use Only

